




Applied geoscience for our
changing Earth

The solid phase distribution and bioaccessibility of potentially harmful elements in natural ironstone soils in the UK

Joanna Wragg

© NERC All rights reserved

Study location



- 7 sampling sites
 - Residential and allotments at each
- Underlying parent geology – Northampton Sand Formation (NSF)
 - Jurassic Ironstone
 - Stratified by
 - Great Oolite
 - Inferior Oolite
 - Lias Group



Study Aim

- Assess the potential availability of As, Cr and Ni in ironstone derived soils from the Cherwell district via the oral ingestion route.
- **Why?**
- The Advanced soil geochemical atlas of England and Wales'

	Min	Max	Mean
As mg kg ⁻¹	0	820	20
Cr mg kg ⁻¹	5.1	1141	68
Ni mg kg ⁻¹	0.26	459	23
- Other studies of Jurassic ironstones have shown soil PHE concentrations above UK soil guideline and generic assessment criteria (at the time of the study)

Land use	Soil Guideline Value (mg kg ⁻¹ DW)		
	Inorganic As	Cr*	Ni
Residential	20 (32)	200 (627)*	75 (130)*
Allotment	20 (43)	130 (15300)	50 (230)


- Most work in the UK has been carried out on As and Pb



© NERC All rights reserved

Study Aim II


- Identification of the source of the bioaccessible PHEs
 - The source components
- Identification of any relationships with the underlying parent geology



© NERC All rights reserved

Experimental

- **Total As, Cr and Ni**
 - 12 residential, 18 allotment
 - Bioaccessible As, Cr and Ni
 - Sieved as per BGS standard geochemical field survey
 - Dried at 35 ± 2°C
 - Sieved to <250 µm




- Auger soil sampling
- Sieving Ø < 250µm



© NERC All rights reserved

BGS Modified PBET

- Study used the BGS Modified PBET
 - Carried out before the UBM was developed enough for use
- Mimics kinetic and chemicals changes in the human GI environment
- Considers Stomach and Small Intestine Conditions
 - 2-3 year old child
 - 37°C
 - Stomach pH @ 2.5
 - Small Intestine pH @ 7.0




© NERC All rights reserved

How PHEs are distributed between the soil components?


CISED Test Chemometric Identification of Substrates and Element Distributions

- Separate aliquots of aqua regia of increasing concentration.
- Passed through the sample under centrifugal force.
- Determination by ICP-AES.
- Chemometric data processing.
- Identification of physico-chemical hosts and the metal distributions within the sample under test.




© NERC All rights reserved

Results – Total PHEs



© NERC All rights reserved


Results – Total PHEs



© NERC All rights reserved

- Total PHE concentrations significantly higher in the LG
- Median LG Ni below new SGV for residential and allotment sites
- Median LG Cr and As below new SGV for residential sites


Results - Bioaccessibility



© NERC All rights reserved

- All results based on highest of the two phases
- All results below the SGV/GAC – current or previous
- Not statistically different (A-R)
 - In general a wider range of values for allotment soils
- Why???


Results - Bioaccessibility



© NERC All rights reserved

- Bioaccessibility significantly higher in soils from the Lias group
- LG bioaccessibility ca. double IO


Results - Bioaccessible Fraction (%)




© NERC All rights reserved

- BAF is the value used in a risk assessment
- % BAF for the IO group is higher for all PHE than LG
 - Opposite to the observations using the actual mg kg⁻¹ bioaccessibility values
- WHY???
- Because of the lower total PHE concentrations for this group


Results - CISED



PHE distributions




Bioaccessible As source components




- Similar for both LG samples
 - Sequential dissolution of organic, fertiliser, carbonate, Mn and Al oxides
 - More As association with Al oxides at Drayton
- Some bioaccessible As with Fe oxide – but this is the host of the non-bioaccessible form
- At other sites – carbonate/high carbonate play a larger host role

Bioaccessible Cr source components



- As with As, sequential dissolution of organic, fertiliser, carbonate, Mn and Al oxides
- Some bioaccessible Cr with Fe oxide – but this is the host of the non-bioaccessible form


Bioaccessible Ni source components



- Geochemical hosts of bioaccessible Ni
 - Organic, fertiliser, carbonate, Mn oxide (small amounts)
- Non bioaccessible hosts
 - High carbonate, Mn, Al and Fe oxides

Conclusions

- The combination of methodologies has been useful for understanding process control
 - Identifying host soil components *etc.*
- Bioaccessible As and Cr appear to result from the same geochemical host components
- Bioaccessible Ni results from more environmentally mobile host components – e.g. carbonates
- Parent geology is key
 - IO PHEs have higher bioaccessibility as a %
- Land use may play an important role in understanding contaminant bioaccessibility
 - Allotments indicate a broader range of bioaccessibility
 - different / changed distributions
 - Additives – fertilisers, manure *etc.*



7th International Workshop on
Chemical Bioavailability in the
Terrestrial Environment

November 2013

Hosted at the

British Geological Survey, Keyworth, Nottingham,
UK

Cbio7@bgs.ac.uk

